

ACAA2 Recombinant antibody

Cat:B35011D

Company: HaoKebio

Uniprot ID:P42765

Applications: IHC:1:100-1:200

Organism:Rabbit

IHC-Polymer:1:400-1:800

Species reactivity:Human Mouse Rat

IHC-TSA:1:500-1:1000

Molecular Weight Calculation: 42 kDa

WB:1:1000-1:5000

Observed Molecular Weight: 42 kDa

Background:

The encoded protein catalyzes the last step of the mitochondrial fatty acid beta-oxidation spiral. Unlike most mitochondrial matrix proteins, it contains a non-cleavable amino-terminal targeting signal.

Protein full name:

Acetyl-CoA Acyltransferase 2

Synonyms:

T1; DSAEC

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Cytoplasm

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:


Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat

at Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Dilution of 1:2500 incubated at room temperature for 1.5 hours.

Source of Reagents:

发表[中文论文]请标注:ACAA2(B35011D)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:ACAA2(B35011D) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.