

IgD Recombinant antibody

Cat:B32104R

Company: Haokebio

Uniprot ID:P01880

Applications: IHC:1:100-1:200

Organism:Rabbit

IHC-Polymer:1:400-1:800

Species reactivity:Human


TSA:1:500-1:1000

Background:

IgD antibody can react with the delta chain of immunoglobulin IgD, and it is used for the differential diagnosis of lymphoma, plasmacytoma, and B-cell-derived Hodgkin lymphoma.

dium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Sample: Human tonsil tissue, 4% PFA 12-24h

Protein full name:

immunoglobulin heavy constant delta

Synonyms:

IGHD

Immunogen:

Peptide

Isotype:

IgG

Subcellular location:

Cytoplasm

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0), Medium high heat for 8 minutes, stop for 7 minutes, medium

Source of Reagents:

发表[中文论文]请标注:IgD(B32104R)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:IgD(B32104R) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.